


The SpinQuest Experiment (E1039) at Fermilab


Ishara Fernando
For the SpinQuest Collaboration



5th Workshop on the
QCD Structure of the Nucleon
Alcalá de Henares, Madrid, SPAIN
October 4-8, 2021



Universidad
de Alcalá



U.S. DEPARTMENT OF
ENERGY

Office of
Science

Outline

- Physics motivation
- Possible missing spin contributions
- TMD PDFs, Sivers Function & Sign
- Global analyses, global context & sea-quark Sivers functions
- Polarized fixed target Drell-Yan / SpinQuest / E1039 experiment at Fermilab
- Projected Uncertainties & goodness of event-reconstruction
- SpinQuest / E1039 timeline
- SpinQuest / E1039 Goals

Physics Motivation

Ji's decomposition

$$\frac{1}{2} = \boxed{\frac{1}{2} \sum_q \Delta q} + \sum_q L_q^z + J_g^z$$

Jaffe-Manohar decomposition

$$\frac{1}{2} = \boxed{\frac{1}{2} \sum_q \Delta q} + \sum_q \mathcal{L}^q + \Delta G + \mathcal{L}^g$$

Intrinsic spin contribution
by valence & sea quarks

$\sim 12\%$

QCD Corrected
Quark Parton Model
(Ellis-Jaffe Sum rule)

0.189 ± 0.005

$$A = \frac{d\sigma^{\uparrow\downarrow} - d\sigma^{\uparrow\uparrow}}{d\sigma^{\uparrow\downarrow} + d\sigma^{\uparrow\uparrow}}$$

$$\int_0^1 g_1^p dx = \boxed{0.126} \pm 0.010 \pm 0.015$$


$$g_1(x) = \frac{1}{2} \sum e_i^2 (q_i^+(x) - q_i^-(x))$$

EMC Collaboration (1989)


Nuclear Physics B328 (1989) 1-35

Asymmetry measurements from Deep inelastic scattering of longitudinally polarized muons on longitudinally polarized proton

Physics Motivation



Nuclear Physics B328 (1989) 1–35



$$\int_0^1 g_1^p dx = 0.126 \pm 0.010 \pm 0.015$$


$$\langle S_z \rangle_{\text{valence}} = +0.535 \pm 0.032 \pm 0.046$$

$$\langle S_z \rangle_{\text{sea}} = -0.475 \pm 0.080 \pm 0.115$$

Intrinsic spin contribution
(total) by valence & sea quarks


$\sim 12\%$

COMPASS Collaboration: Physics Letters B 753 (2016) 18–28



Possible missing spin contributions

K.-F. Liu et al
arXiv:1203.6388



$\Delta\Sigma_q \approx 25\%$
 $2 L_q \approx 50\%$ (4% (valence)+46% (sea))
 $2 J_g \approx 25\%$

$$\frac{1}{2} = \frac{1}{2} \Delta\Sigma + \boxed{\Delta G + L_g} + \boxed{L_q} + \boxed{L_{\bar{q}}}$$

Gluon total angular momentum


Valence quarks' OAM

Sea-quarks' OAM


Jaffe-Manohar decomposition

- Sea quark OAM could be a major contribution (J. Ellis and M. Karliner, Phys. Lett. B213 (1988) 73)
- Separation of gluon intrinsic spin and OAM is constrained by gauge invariance

C. Alexandrou et al
PRL 119, 142002 (2017)



	$\frac{1}{2} \Delta\Sigma$	J	L	$\langle x \rangle$
u	0.415(13)(2)	0.308(30)(24)	-0.107(32)(24)	0.453(57)(48)
d	-0.193(8)(3)	0.054(29)(24)	0.247(30)(24)	0.259(57)(47)
s	-0.021(5)(1)	0.046(21)(0)	0.067(21)(1)	0.092(41)(0)
g	-	0.133(11)(14)	-	0.267(22)(27)
tot.	0.201(17)(5)	0.541(62)(49)	0.207(64)(45)	1.07(12)(10)



TMD PDFs

		Quark Polarization		
		U	L	T
Nucleon Polarization	U	$f_1 = \odot$	N/A	$h_1^\perp = \odot - \ominus$ Boer-Mulders
	L	N/A	$g_{1L} = \odot - \ominus$ Helicity	$h_{1L}^\perp = \odot - \ominus$
	T	$f_{1T}^\perp = \odot - \ominus$ Sivers	$g_{1T}^\perp = \odot - \ominus$	$h_1 = \odot - \ominus$ $h_{1T}^\perp = \odot - \ominus$ Transversity

$$\Phi(x, k_T; S) = \int \frac{d\xi^- d\xi_T}{(2\pi)^3} e^{ik \cdot \xi} \langle P, S | \bar{\psi}(0) \mathcal{U}_{[0, \xi]} \psi(\xi) | P, S \rangle |_{\xi^+ = 0}$$




Quark correlator can be decomposed into 8 components
(6 T-even and 2 T-odd terms) at leading-twist

$$\begin{aligned} \Phi(x, k_T, P, S) = & f_1(x, k_T^2) \frac{\not{P}}{2} + \frac{h_{1T}(x, k_T^2)}{4} \gamma_5 [\not{S}_T, \not{P}] + \frac{S_L}{2} g_{1L}(x, k_T^2) \gamma_5 \not{P} + \frac{k_T \cdot S_T}{2M} g_{1T}(x, k_T^2) \gamma_5 \not{P} \\ & + S_L h_{1L}^\perp(x, k_T^2) \gamma_5 \frac{[k_T, \not{P}]}{4M} + \frac{k_T \cdot S_T}{2M} h_{1T}^\perp(x, k_T^2) \gamma_5 \frac{[k_T, \not{P}]}{4M} \end{aligned}$$

T-even


$$+ ih_1^\perp(x, k_T^2) \frac{[k_T, \not{P}]}{4M} - \frac{\epsilon_T^{k_T S_T}}{4M} f_{1T}^\perp(x, k_T^2) \not{P}$$

T-odd



TMD PDFs

Polarized Semi-Inclusive DIS



SIDIS

$$\frac{d\sigma_{SIDIS}^{LO}}{dx dy dz dp_T^2 d\phi_h d\psi} = \left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\epsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] \times (F_{UU,T} + \epsilon F_{UU,L}) \left\{ 1 + \cos 2\phi_h (\epsilon A_{UU}^{\cos 2\phi_h}) \right. \\ \left. + S_T \begin{bmatrix} \sin(\phi_h - \phi_S) (A_{UT}^{\sin(\phi_h - \phi_S)}) \\ + \sin(\phi_h + \phi_S) (\epsilon A_{UT}^{\sin(\phi_h + \phi_S)}) \\ + \sin(3\phi_h - \phi_S) (\epsilon A_{UT}^{\sin(3\phi_h - \phi_S)}) \end{bmatrix} \right\}$$

PDF \otimes FF

$$A_{UU}^{\cos 2\phi_h} \propto h_1^{\perp q} \otimes H_{1q}^{\perp h}$$


$$A_{UT}^{\sin(\phi_h - \phi_S)} \propto f_{1T}^{\perp q} \otimes D_{1q}^h$$

$$A_{UT}^{\sin(\phi_h + \phi_S)} \propto h_1^q \otimes H_{1q}^{\perp h}$$

$$A_{UT}^{\sin(3\phi_h - \phi_S)} \propto h_{1T}^{\perp q} \otimes H_{1q}^{\perp h}$$

BM \otimes CF
Sivers \otimes FF
Transv \otimes CF
Pretz \otimes CF

Polarized Drell-Yan



DY

$$\frac{d\sigma^{LO}}{d\Omega} = \frac{\alpha_{em}^2}{Fq^2} F_U^1 \left\{ 1 + \cos^2 \theta + \sin^2 \theta \cos 2\varphi_{CS} A_U^{\cos 2\varphi_{CS}} \right. \\ \left. + S_T \begin{bmatrix} (1 + \cos^2 \theta) \sin \varphi_S A_T^{\sin \varphi_S} \\ + \sin^2 \theta \left(\sin(2\varphi_{CS} + \varphi_S) A_T^{\sin(2\varphi_{CS} + \varphi_S)} \right. \right. \\ \left. \left. + \sin(2\varphi_{CS} - \varphi_S) A_T^{\sin(2\varphi_{CS} - \varphi_S)} \right) \right\}$$

beam target

PDF \otimes PDF

BM \otimes BM
 f_1 \otimes Sivers
BM \otimes Transv
BM \otimes Pretz

$$A_T^{\cos 2\varphi_{CS}} \propto h_1^{\perp q} \otimes h_1^{\perp q}$$

$$A_T^{\sin \varphi_S} \propto f_1^q \otimes f_{1T}^{\perp q}$$

$$A_T^{\sin(2\varphi_{CS} - \varphi_S)} \propto h_1^{\perp q} \otimes h_{1T}^{\perp q}$$

$$A_T^{\sin(2\varphi_{CS} + \varphi_S)} \propto h_1^{\perp q} \otimes h_1^q$$

$$h_1^{\perp q} \Big|_{SIDIS} = -h_1^{\perp q} \Big|_{DY}$$

$$f_{1T}^{\perp q} \Big|_{SIDIS} = -f_{1T}^{\perp q} \Big|_{DY}$$

$$h_1^q \Big|_{SIDIS} = h_1^q \Big|_{DY}$$


$$h_{1T}^{\perp q} \Big|_{SIDIS} = h_{1T}^{\perp q} \Big|_{DY}$$

* For these two processes TMD factorization is proven




Sivers Function

$$f_{q/p^\uparrow}(x, \mathbf{k}_T) = f_{q/p}(x, \mathbf{k}_T) + f_{1T}^\perp(x, \mathbf{k}_T) \mathbf{S} \cdot (\hat{\mathbf{P}} \times \hat{\mathbf{k}}_T)$$

The Sivers function describes the correlation between the momentum direction of the struck quark and the spin of its parent nucleon.





- The gauge-invariant definition of the Sivers function predicts the opposite sign for the Sivers function in SIDIS compared to processes with color charges in the initial state and a colorless final state in Drell-Yan, J/ψ , W^\pm , Z
- This inclusion of the gauge link has profound consequences on factorization proofs and on the concept of universality, which are of fundamental relevance for high-energy hadronic physics




Sign of Sivers Functions

STAR Collaboration (PRL 116 132301 (2016))



COMPASS Collaboration (PRL 119 112002 (2017))




TSSA amplitude for W^+/W^- from STAR data is favors the “sign-change”
 In DY relative to SIDIS (model based without TMD evolution)

Dark Shaded (Light-shaded): with(without)
 “sign-change”


Global analyses: Sivers functions

A. Bacchetta, F. Delcarro,
C. Pasiano, M. Radici
arXiv 2004.14278 (2020)




HERMES (2020)
COMPASS (2009)
COMPASS (2015)
JLab (2011)


M. Anselmino, M. Boglion, U. D'Alesio, S. Melis, F. Murgia, A. Prokudin_PRD 79_54010_(2009)



HERMES (2020), COMPASS (2009), COMPASS (2015)
JLab (2011), STAR (2016), COMPASS DY (2017)






M. Bury, A. Prokudin, A. Vladimirov, JHEP_05_151 (2021)



HERMES (2009)
COMPASS (2009)
COMPASS (2015)
JLab (2011)
STAR (2016)

M. Echevarria, Z. Kang, J. Terry_JHEP_01_126_(2021)

SpinQuest in the Global Context





- The E1039 experiment will provide unique information for the sea quark Sivers function in DY
- ←Plot: Uncertainties in the predicted Sivers asymmetry in polarized Drell-Yan reactions. (Negative x_F is dominated by valence quarks, while positive x_F is dominated by sea quarks)

Drell-Yan measurements above the J/ψ peak fall in a unique region with Q^2 in the range of $16 < M^2 < 81$ GeV² and $Q_T < \text{few GeV}$

Sea-quarks Sivers functions

STAR Collaboration (PRL 116 132301 (2016))




The solid gray bands represent the uncertainty due to the unknown sea quark Sivers functions estimated by saturating the sea quark Sivers function to their positivity limit in the KQ (Z.-B. Kang and J. -W. Qiu PRL 103,172001 (2009)) calculation

- Initial attempts to measure the Sivers asymmetry for sea quark Sivers have been reported by the STAR collaboration at RHIC using W/Z boson production. Their data is statistically limited and favor a sign-change only if TMD evolutions effects are significantly smaller than expected.
- SpinQuest will perform the first measurement of the Sivers asymmetry in Drell-Yan proton-proton scattering from the sea quarks.

Polarized fixed target Drell-Yan : Sensitivity to sea-quarks

beam: valence quarks
at high x

target: sea quarks at
low/intermediate x




Sea-quarks
dominance

$$\frac{d^2\sigma}{dx_b dx_t} = \frac{4\pi\alpha^2}{x_b x_t S} \sum_{q \in \{u, d, s, \dots\}} e_q^2 [\bar{q}_t(x_t) q_b(x_b) + \cancel{q_t(x_t) \bar{q}_b(x_b)}]$$

u-quark dominance
(2/3)² vs. (1/3)²

acceptance limited
(Fixed Target, Hadron Beam)



Valence-quarks
dominance

Polarized fixed target DY & J/ψ @ SpinQuest / E1039 experiment


$$A = \frac{\sigma(p_b^{un} p_t^\uparrow) - \sigma(p_b^{un} p_t^\downarrow)}{\sigma(p_b^{un} p_t^\uparrow) + \sigma(p_b^{un} p_t^\downarrow)}$$

Drell-Yan $\sigma(p + p^{\uparrow(\downarrow)} \rightarrow \gamma + X)$

$$f_{q/p^\uparrow}(x, \mathbf{k}_T, \mathbf{S}_T; Q) = f_{q/p}(x, \mathbf{k}_T; Q) + \frac{1}{2} \Delta^N f_{q/p^\uparrow}(x, \mathbf{k}_T, \mathbf{S}_T; Q)$$

J/ψ $\sigma(p + p^{\uparrow(\downarrow)} \rightarrow J/\psi + X)$

$$f_{g/p^\uparrow}(x, \mathbf{k}_T, \mathbf{S}_T; Q) = f_{g/p}(x, \mathbf{k}_T; Q) + \frac{1}{2} \Delta^N f_{g/p^\uparrow}(x, \mathbf{k}_T, \mathbf{S}_T; Q)$$



- SpinQuest will be able to explore a new region of kinematics for J/ψ compare to the PHENIX measurements
- J/ψ production:
 - PHENIX $\rightarrow gg$ fusion at $\sqrt{s} = 200$
 - SpinQuest $\rightarrow q\bar{q}$ annihilation at $\sqrt{s} = 15.5$

About SpinQuest/E1039 Collaboration

<https://spinquest.fnal.gov>

INSTITUTION **20**

FULL MEMBERS **53** Postdocs **7** Grad. Students **15**

AFFILIATE MEMBERS

[1\) Abilene Christian University](#)

Donald Isenhower (PI), Michael Daugherty, Shon Watson

Haley Stien, John Marsden, Mitchell Schneller, Nathan Rowlands, Roy Salinas, Rusty Towell, Shannon McNease, Yves Ngenzi, Thomas Fitch

[2\) Argonne National Laboratory](#)

Paul Reimer (PI), Donald Geesaman

Kevin Bailey, Thomas O'Connor

[3\) Aligarh Muslim University](#)

Huma Haider (PI)

[4\) Boston University](#)

David Sperka (PI), Zijie Wan

[5\) Fermi National Accelerator Laboratory](#)

Richard Tesarek (PI), Carol Johnstone, Charles Brown

[6\) KEK](#)

Shin'ya Sawada (PI)

[7\) Los Alamos National Laboratory](#)

Kun Liu (SP), Ming Liu, Astrid Morreale, Mikhail Yurov, Kei Nagai, Zongwei Zhang

[8\) Mississippi State University](#)

Lamiaa El Fassi (PI), Catherine Ayuso, Nuwan Chaminda

[9\) New Mexico State University](#)

Stephen Pate (PI), Vassili Papavassiliou, Abinash Pun, Forhad Hossain, Dinupa Nowarathne

[10\) RIKEN](#)

Yuji Goto (PI)

[11\) Shandong University](#)

Qinghua Xu (PI), Zhaohuizi Ji

[12\) Tokyo Institute of Technology](#)

Toshi-Aki Shibata (PI)

[13\) University of Colombo](#)

Darshana Perera (PI), Harsha Sirilal, Vibodha Bandara

[14\) University of Illinois, Urbana-Champaign](#)

Jen-Chieh Peng (PI), Ching Him Leung

[15\) University of Michigan](#)

Wolfgang Lorenzon (PI), Levgen Lavrukhin, Noah Wuerfel

[16\) University of New Hampshire](#)

Karl Slifer (PI), David Ruth

[17\) Tsinghua University](#)

Zhihong Ye (PI)

[18\) University of Virginia](#)

Dustin Keller (SP), Kenichi Nakano, Ishara Fernando, Zulkaida Akbar, Waqar Ahmed, Liliet Diaz, Anchit Arora, Arthur Conover, Jay Roberts

Shigeru Ishimoto

Jan Boissevain, Melynda Brooks, Matt Durham, David Kleinjan, Sho Uemura, Cesar Da Silva, Patrick McGaughey, Andi Klein

Dipangkar Dutta

Naomi Makins, Daniel Jumper, Jason Dove, Mingyan Tian, Bryan Dannowitz, Randall McClellan, Shivangi Prasad

Daniel Morton, Richard Raymond, Marshall Scott

Maurik Holtrop

Donal Day, Donald Crabb, Jixie Zhang, Oscar Rondon, Arthur Conover, Brandon Kriesten, Simonetta Liuti, Ellen Brown, Blaine Norum, Matthew Roberts

Takahiro Iwata, Norihiro Doshita



[19\) Yamagata University](#)

Yoshiyuki Miyachi (PI), Norihito Doshita

[20\) Yerevan Physics Institute](#)

Hrachya Marukyan (PI)

Fermilab proton beam main injector



- 120 GeV/c proton beam
- $\sqrt{s} = 15.5$ GeV
- Projected beam
 - ❖ 5×10^{12} protons/spill Where $spill \approx 4.4$ s/min
 - ❖ Bunches of 1ns with 19ns intervals ~ 53 MHz
 - ❖ 7×10^{17} protons/year on target!

Fermilab proton beam main injector

$$\frac{d^2\sigma}{dx_1 dx_2} = \frac{4\pi\alpha^2}{9x_1 x_2} \frac{1}{s} \times \sum_i e_i^2 [q_{ti}(x_t)\bar{q}_{bi}(x_b) + \bar{q}_{ti}(x_t)q_{bi}(x_b)]$$

Fermilab E866/NuSea

Data in 1996-1997

^1H , ^2H and nuclear targets

800 GeV proton beam

Fermilab E906/E1039


Data in > 2010

^1H , ^2H and nuclear targets



120 GeV proton beam

Therefore, the SpinQuest/E1039 experiment will get,

- Cross-Section scales as **~7** times compare to that with 800 GeV beam
- Luminosity is **~7** times compare to that with 800 GeV beam
- **~49** x Statistics with 800 GeV beam



SpinQuest / E1039 Experiment Setup




Solid Iron Focusing Magnet, Hadron absorber and beam dump

Station 1: Hodoscope array MWPC tracking

Station 2 and 3: Hodoscope array Drift Chamber tracking


Station 4: Hodoscope array Prop tube tracking



Drawing: T. O'Connor and K. Bailey

Polarized solid NH₃ & ND₃ target setup

- ❖ Designed for high intensity proton beam (5×10^{12} protons/spill with 4.4s spill) by LANL-UVA group
- ❖ 8 cm long solid NH₃ and ND₃ target cells
- ❖ Magnetic Field: $B = 5$ T with uniformity $dB/B < 10^{-4}$ over 8 cm
- ❖ ⁴He evaporation refrigerator (3 W of maximum cooling power) keeping the target at 1.1 K.
- ❖ 140 GHz microwave source (with DNP technique)




Predicted Uncertainties

- Beam ($\sim 2.5\%$)
 - Relative luminosity ($\sim 1\%$)
 - Drifts ($< 2\%$)
 - Scraping ($\sim 1\%$)

- Analysis sources ($\sim 3.5\%$)
 - Tracking efficiency ($\sim 1.5\%$)
 - Trigger & geometrical acceptance ($< 2\%$)
 - Mixed background ($\sim 3\%$)
 - Shape of DY ($\sim 1\%$)

- Target ($\sim 6-7\%$)
 - TE calibration (proton $\sim 2.5\%$; deuteron $\sim 4.5\%$)
 - Polarization inhomogeneity ($\sim 2\%$)
 - Density of target ($\text{NH}_3(\text{s})$) ($\sim 1\%$)
 - Uneven radiation damage ($\sim 3\%$)
 - Beam-Target misalignment ($\sim 0.5\%$)
 - Packing fraction ($\sim 2\%$)
 - Dilution factor ($\sim 3\%$)





DGLAP: M. Anselmino et al arXiv:1612.06413
 TMD-1: M. G. Echevarria et al arXiv:1401.5078
 TMD-2: P. Sun and F. Yuan arXiv:1308.5003
 A. Prokudin et al (in progress)
 I. Fernando, D. Keller (in progress)

$$A = \frac{2}{f|S_T|} \frac{\int d\phi_S d\phi \frac{dN(x_b, x_t, \phi_S, \phi)}{d\phi_S d\phi} \sin(\phi_S)}{N(x_b, x_t)}$$

x_2 bin	$\langle x_2 \rangle$	$\text{NH}_3 (p^\uparrow)$		$\text{ND}_3 (d^\uparrow)$	
		N	ΔA (%)	N	ΔA (%)
0.10 - 0.16	0.139	5.0×10^4	3.2	5.8×10^4	4.3
0.16 - 0.19	0.175	4.5×10^4	3.3	5.2×10^4	4.6
0.19 - 0.24	0.213	5.7×10^4	2.9	6.6×10^4	4.1
0.24 - 0.60	0.295	5.5×10^4	3.0	6.4×10^4	4.1

Material	Density	Dilution factor	Packing fraction	Polarization	Interaction length
NH_3	0.867 g/cm^3	0.176	0.60	80%	5.3%
ND_3	1.007 g/cm^3	0.300	0.60	32%	5.7%

Goodness of event-reconstruction from E906



- Monte-Carlo describe data well
- Better resolution than expected

- $\delta\sigma_M(J/\psi) \sim 220 \text{ MeV}$
- $\delta\sigma_M(DY) \sim$ truth-reconstructed from event-by-event MC
- J/ψ and ψ' separation

The projected event selection/reconstruction is expected to be the same for E1039

SpinQuest / E1039 Timeline

- 2018, March: DOE approval
- 2018, May: Fermilab stage-2 approval
- 2018, June: E906 decommissioned
- 2019, May: Transferred the polarized target from UVA to Fermilab
- Now: commission all components using cosmic rays
- Polarized target to be installed by Spring of 2022
- E1039 commissioning starts in the beginning of 2022
[Run for 2+ years, 2022-2024+]

SpinQuest / E1039 Goals

- SpinQuest will perform the first measurement of the Sivers asymmetry in Drell-Yan proton-proton scattering from the sea quarks (\bar{u} & \bar{d}) with sign.

$$f_{1T}^\perp|_{\text{SIDIS}} = - f_{1T}^\perp|_{\text{DY}}$$

A direct QCD prediction is a Sivers effect in the Drell-Yan process that has the opposite sign compared to the one in semi-inclusive DIS.

- Measurement of Sivers function for gluons (J/psi TSSA)
- Explore a unique range of virtualities and transverse momenta not accessible through Z^0/W^\pm measurements
- Extensions: transversity, tensor charge, tensor polarized observables, dark sector, polarized proton beam,...

Welcome!


Please Join The Effort

Dustin Keller (dustin@virginia.edu)[Spokesperson]

Kun Liu ([Spokesperson])

<https://spinqest.fnal.gov/>

<http://twist.phys.virginia.edu/E1039/>



Thank you



UNIVERSITY
of
VIRGINIA



U.S. DEPARTMENT OF
ENERGY

Office of
Science

This work is supported by DOE contract DE-FG02-96ER40950

Back-up Slides

(Un)Polarized DY

Experiment	Particles	Energy (GeV)	x_b or x_t	Luminosity ($\text{cm}^{-2} \text{s}^{-1}$)	P_b or P_t (f)	rFOM [#]	Timeline
COMPASS (CERN)	$\pi^- + p^\uparrow$	160 GeV $\sqrt{s} = 17$	$x_t = 0.1 - 0.3$	2×10^{33}	$P_t = 90\%$ $f = 0.22$	1.1×10^{-3}	2015-2016, 2018
J-PARC (high-p beam line)	$\pi^- + p$	10-20 GeV $\sqrt{s} = 4.4-6.2$	$x_b = 0.2 - 0.97$ $x_t = 0.06 - 0.6$	2×10^{31}	---	---	>2020? under discussion
fsPHENIX (RHIC)	$p^\uparrow + p^\uparrow$	$\sqrt{s} = 200$ $\sqrt{s} = 510$	$x_b = 0.1 - 0.5$ $x_b = 0.05 - 0.6$	8×10^{31} 6×10^{32}	$P_b = 60\%$ $P_b = 50\%$	4.0×10^{-4} 2.1×10^{-3}	>2021?
SeaQuest (FNAL: E-906)	$p + p$	120 GeV $\sqrt{s} = 15$	$x_b = 0.35 - 0.9$ $x_t = 0.1 - 0.45$	3.4×10^{35}	---	---	2012 - 2017
Pol tgt DY[‡] (FNAL: E-1039)	$p + p^\uparrow$ $p + d^\uparrow$	120 GeV $\sqrt{s} = 15$	$x_t = 0.1 - 0.45$	3.0×10^{35} 3.5×10^{35}	$P_t = 85\%$ $f = 0.176$	0.15	2021-2023+
Pol beam DY[§] (FNAL: E-1027)	$p^\uparrow + p$	120 GeV $\sqrt{s} = 15$	$x_b = 0.35 - 0.9$	2×10^{35}	$P_b = 60\%$	1	>2021?

Source: Wolfgang Lorenzon

[‡] 8 cm NH₃ target / [§] $L = 1 \times 10^{36} \text{ cm}^{-2} \text{ s}^{-1}$ (LH₂ tgt limited) / $L = 2 \times 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$ (10% of MI beam limited)

*not constrained by SIDIS data / [#]rFOM = relative lumi * P² * f² wrt E-1027 (f=1 for pol p beams, f=0.22 for π^- beam on NH₃)